

## APPLICATION

For control of aphid and greenhouse whitefly, on ornamental, tomato and eggplant crops.

For control of caterpillar pests in apple and kiwifruit crops

Use at a rate of 10 g per litre of spray, or as specified below .

### Specific Instructions for Use

Add to warm water and mix well first to dissolve, then add to spray tank last to avoid excessive foaming.

Maintain agitation of spray mixture at all times.

Ensure adequate coverage of all plant surfaces, particularly undersides of lower leaves.

### Application Rates

#### For Greenhouse Whitefly Control

##### High Population Pressure and Existing Infestation

For existing infestations apply at 3 day intervals for 2 weeks. Then resume regular weekly sprays.

##### Moderate to Low Population Pressure and Existing Infestation

Apply at 7 day intervals for at least 4 applications. Monitor pest nymph numbers on a weekly basis.

##### Low Population Pressure - no infestation

Apply at 5g per litre. Repeat Application at 2 weekly intervals. Monitor Yellow traps for early detection of pest adults

USEAGE TABLE		Boom Spray	Motorised Knapsack or Trolley HandGun or Knapsack	MistBlower
Crops	Pest			
Ornamentals	Greenhouse Whitefly, Aphid			
Tomato	Product Rate per litre	10g (1kg/100 l)	10g (100g per 10 litres)	40 g (600g per 15 litre)
Eggplant	Water Rate per 1000m <sup>2</sup>	200 litre	250 litre	50 litre
Conventional Orchard Airblast Sprayer				
Kiwifruit	Leafroller	Apply 20kg in 2000 litres water per hectare Apply from Mid February to 14 days before harvest at 21 day intervals		
Apple	Leafroller, Codlin Moth	Apply 20 kg in 2000 litres water per hectare. Apply at 14 day intervals during the season commencing at petal fall		

NOT Recommended for use in LVM, CDA or Fog applicators. NOT for use in irrigation, fertigation, or nutrient solution.

NOT for application to soil, water or any growing media

### Storage Handling and Compatibility

**Store product dry and sealed under dark cool conditions.**

Use the spray within 8 hours of mixing. Apply towards early evening.

Applicators should wear eye and skin protection. Respirator should be worn while measuring powder.

Empty contents of sprayer and wash tank and nozzles after use

Do NOT mix with other spray products unless prior experience shows the combination to be safe and effective. Additional wetting /sticking agents are not usually required.

No Re-entry restriction. Wash exposed skin after application. Wash residues off produce prior to sale.

Manufacturing Date  
Batch Number

Quality Check

### Shelf Life

When stored appropriately this product shows no significant degradation for 2 Years from the date of manufacture. Contact your supplier for further information on the use of any product that is older than this.

Registered pursuant to the ACVM Act 1997 REGISTRATION NUMBER P 7363  
see [www.nzfsa.govt.nz/acvm](http://www.nzfsa.govt.nz/acvm) for registration conditions.

Manufacturer Suntec (NZ) Ltd .P O Box 41 Tokomaru 5450. New Zealand . +64 6 3298142 . Fax +64 6 3298041 .Email [suntec@suntec.co.nz](mailto:suntec@suntec.co.nz)

Contents in sealed 2.5 kg packages